

Strange baryon production asymmetry in $K^\pm N$ interactions

G. H. Arakelyan¹, C. Merino², and Yu. M. Shabelski^{3a}

¹ YerPhI, Armenia

² Dpto. de Física de Partículas, Facultade de Física, and IGAE, Universidade de Santiago de Compostela, Galicia, Spain

³ SPNPI, Gatchina, St.Petersburg, Russia

Received: date / Revised version: date

Abstract. The asymmetry of strange baryon production in Kp interactions at high energies is considered in the framework of the Quark-Gluon String Model. The contribution of the string-junction mechanism to the strange baryon production is analysed.

PACS. 12.40.-y Other models for strong interactions – 13.60.Rj Baryon production – 13.75.Jz Kaon-baryon interactions

The present report is devoted to the calculation of asymmetry of strange baryon production in the case of kaon beams and to the analysis of the contribution of the string-junction (SJ) mechanism in Kp collisions.

It is very important to understand the role of the SJ mechanism in the dynamics of high-energy hadronic interactions, in particular in processes implying baryon number transfer. Significant results on this question were obtained in [1, 2, 3, 4], where the SJ mechanism was used to analyse the strange baryon production in πp and pp interactions. The detailed analysis of SJ contribution into strange baryon production in KN interaction was presented in [5]

In the present paper we analyse the existing data on asymmetry of Λ and $\bar{\Lambda}$ production on K -beams [6].

We compare the experimental data with the result of our calculations for a value of the SJ intercept $\alpha_{SJ} = 0.9$.

The formulas describing the inclusive spectrum (i.e., Feynman- x , x_F , distribution) of a secondary hadron h in KN scattering in the Quark Gluon String Model (QGSM) frame were presented in papers [5, 7].

The complete set of distribution and fragmentation functions used in this paper is presented in [5].

The SJ mechanism has a nonperturbative origin and since it is at present not possible to determine the value of α_{SJ} in QCD from first principles. Thus we treat α_{SJ} and ε (the weight of the diagram describing the SJ contribution, see [5]) as phenomenological parameters which should be determined from experimental data. In the present calculation, we use the values $\alpha_{SJ} = 0.9$ and $\varepsilon = 0.024$, as it was done in [2, 5, 8].

As it was shown in [8], the better agreement of QGSM with data on strange baryon production on nucleus was obtained with $\delta = 0.32$, instead of the previous value $\delta = 0.2$. In principle one cannot exclude the possibility that the value of δ would be different for secondary baryons and for mesons (i.e., for Λ -baryon and for kaon).

The fragmentation functions into $\bar{\Lambda}$ do not depend on the SJ mechanism, so the $\bar{\Lambda}$ spectra obtained for different values of α_{SJ} are the same, and they have a very small dependence on the strange quark suppression factor δ (see [5]).

In Fig. 1 we show the comparison of the QGSM calculations with the data on the $\bar{\Lambda}/\Lambda$ asymmetry $A(\bar{\Lambda}/\Lambda)$, produced in K^+p (Fig. 1a) and K^-p (Fig. 1b) interactions at 250 GeV/c [6].

The asymmetry data are rather interesting. In the proton fragmentation region the values of $A(\bar{\Lambda}/\Lambda)$ are close to unity, and that is natural since a proton fragments into Λ with significantly larger probability than into $\bar{\Lambda}$. In the kaon fragmentation region $A(\bar{\Lambda}/\Lambda)$ becomes negative and decreases very fast in the case of K^+ beam at the x_F values where experimental data exist. In the case of K^- beam $A(\bar{\Lambda}/\Lambda)$ increases very fast with x_F and the variation among calculations with different values of the parameters is rather small. Both these behaviors are also natural since the K^+ contains a $\bar{s}$ valence quark which preferably fragments into $\bar{\Lambda}$, while the valence s quark in the K^- fragments rather often into Λ . However, in both cases the $A(\bar{\Lambda}/\Lambda)$ experimental x_F -dependences are much steeper than the theoretical predictions. This is a probable indication that the fragmentation functions $s \rightarrow \Lambda$ and $\bar{s} \rightarrow \bar{\Lambda}$ should be further enhanced.

In the K^+ fragmentation region (Fig. 1a) the predicted values of $A(\bar{\Lambda}/\Lambda)$ at $x_F > 0.4$ show a change of behavior since they start increasing. In this region the contribution of the direct fragmentation of $\bar{s} \rightarrow \bar{\Lambda}$ which makes $A(\bar{\Lambda}/\Lambda)$

^a Contribution to the proceedings of the IV International Conference on Quarks and Nuclear Physics (QNP06), Universidad Complutense, Madrid (Spain), 5-10 June 2006.




Fig. 1. The $\bar{\Lambda}/\Lambda$ asymmetry in (a) K^+p and (b) K^-p collisions. Experimental data at 250 GeV/c [6] and the corresponding QGSM description. The solid curve corresponds to $\alpha_{SJ} = 0.9$, $\varepsilon = 0.024$, and $\delta = 0.32$, the dashed curve to $\delta = 0.2$, and the dashed-dotted curve to $\varepsilon = 0$.

to decrease becomes smaller than the effect of SJ diffusion which increases the multiplicity of Λ . The measurement of the asymmetry $A(\bar{\Lambda}/\Lambda)$ in the region $x_F \geq 0.4$ in K^+p collisions could make the situation more clear.

The predictions for the asymmetry in Ξ and Ω baryon production in K^+p and K^-p interactions are presented in Fig. 2. Here the general situation is similar to that of the case of asymmetry in Λ production shown in Fig. 1. In Fig. 2 we present the predictions for asymmetries in Ξ^- and Ω production in K^+p and K^-p collisions at 250 GeV/c. In the central region of K^+p collisions the yields of Ξ^- and $\bar{\Xi}^+$, as well as those of Ω^- and $\bar{\Omega}^+$ are predicted to be practically the same. The smaller fragmentation function of valence $\bar{s}$ quark into strange baryon is compensated by the larger fragmentation function of the target diquark. In the case of K^- beam (Fig. 2b) the asymmetry for both Ξ and Ω productions increases in the whole region of positive x_F .

The situation for K^- beam seems worse than in the K^+ case since all curves are, as a rule, below the experimental data, but both the number and the quality of experimental data on K^- beam are not very high.

The QGSM predicts a weak energy dependence of the Λ and $\bar{\Lambda}$ production cross-section in Kp collisions at the considered energies.

The experimental data on high-energy Λ production are not in contradiction with the possibility of baryon charge transfer over large rapidity distances, and the $\bar{\Lambda}/\Lambda$ asymmetry is provided by SJ diffusion through baryon charge transfer.




Fig. 2. QGSM prediction for the x_F -dependence of the asymmetry of heavy strange baryons in (a) K^+p and (b) K^-p collisions at 250 GeV/c. Solid curves are for Ξ^- , dashed curves for Ω^- .

The presence of baryon asymmetry in the projectile hemisphere for Kp collisions provides good evidence for such a mechanism.

This work was partially financed by CICYT of Spain through contract FPA2002-01161, and by Xunta de Galicia (Spain) through contract PGIDIT03PXIC20612PN. G. H. Arakelyan and C. Merino were also supported by NATO grant CLG.980335 and Yu. M. Shabelski by grants NATO PDD (CP) PST.CLG 980287 and RCGSS-1124.2003.2. G.H.A. thanks Xunta de Galicia (Spain) for financial support.

References

1. G. H. Arakelyan, A. Capella, A. B. Kaidalov, and Yu. M. Shabelski, Eur. Phys. J. C **26**, 81 (2002).
2. F. Bopp and Yu. M. Shabelski, Yad. Fiz. **68**, 2155 (2005) and hep-ph/0406158 (2004).
3. G. H. Arakelyan, C. Merino, and Yu. M. Shabelski, Yad. Fiz. **69**, N.5 (2006) (in press) and hep-ph/0505100 (2005).
4. O. I. Piskounova, Proc. of the HERA-LHC Workshop, DESY, March 2005; Yad. Fiz. (in press).
5. G. H. Arakelyan, C. Merino, and Yu. M. Shabelski, Yad. Fiz. (in press) and hep-ph/0604103 (2005).
6. G. A. Alves et al. (E769 Collaboration), Phys. Lett. B **559**, 179 (2003) and hep-ex/0303027.
7. Yu. M. Shabelski, Yad. Fiz. **49**, 1081 (1989).
8. F. Bopp and Yu. M. Shabelski, hep-ph/0603193 (2006).